



**QUEEN'S
UNIVERSITY
BELFAST**

Genomic and archaeological evidence suggest a dual origin of domestic dogs

McCormick, F. (2016). Genomic and archaeological evidence suggest a dual origin of domestic dogs. *Science*, 352(6290), 1228-1231. <https://doi.org/10.1126/science.aaf3161>

Published in:
Science

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright the authors 2016.

This is the author's version of the work. It is posted here by permission of the AAAS for personal use, not for redistribution. The definitive version was published in *Science* vol. 352 on 3/06/2016, DOI:10.1126/science.aaf3161.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Title: Genomic and archaeological evidence suggest a dual origin of domestic dogs

Authors: Laurent A. F. Frantz^{1†*}, Victoria E. Mullin^{2†}, Maud Pionnier-Capitan^{3,4}, Ophélie Lebrasseur¹, Morgane Ollivier³, Angela Perri⁵, Anna Linderholm^{1,6}, Valeria Mattiangeli², Matthew D. Teasdale², Evangelos A. Dimopoulos^{1,7}, Anne Tresset⁴, Marilyne Duffraisse³, Finbar McCormick⁸, László Bartosiewicz⁹, Erika Gál¹⁰, Éva A. Nyerges¹⁰, Mikhail V. Sablin¹¹, Stéphanie Bréhard⁴, Marjan Mashkour⁴, Adrian Bălăşescu¹², Benjamin Gillet³, Sandrine Hughes³, Olivier Chassaing³, Christophe Hitte¹³, Jean-Denis Vigne⁴, Keith Dobney¹⁴, Catherine Hänni³, Daniel G. Bradley^{2*} and Greger Larson^{1*}

Affiliations:

¹ The Palaeogenomics & Bio-Archaeology Research Network, Research Laboratory for Archaeology and History of Art, The University of Oxford, Oxford, UK.

² Smurfit Institute of Genetics, Trinity College Dublin, Dublin, Dublin 2, Ireland.

³ CNRS/ENS de Lyon, IGFL, UMR 5242 and French National Platform of Paleogenetics, PALGENE, Ecole Normale Supérieure de Lyon, 46 allée d'Italie, 69364 Lyon Cedex 07, France / Université Grenoble Alpes, Laboratoire d'Ecologie Alpine (LECA), F-38000, Grenoble, France.

⁴ CNRS/MNHN/SUs – UMR 7209, Archéozoologie et Archéobotanique, Sociétés, Pratiques et Environnements, Département Ecologie et Gestion de la Biodiversité, 55 rue Buffon, 75005 Paris, France.

⁵ Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany.

⁶ Department of Anthropology, Texas A&M University, College Station, TX 77843-4352, USA.

⁷ School of Biology, Aristotle University of Thessaloniki, Thessaloniki, Greece.

⁸ School of Geography, Archaeology and Palaeoecology, Queen's University Belfast, University Road, Belfast, Northern Ireland, UK.

⁹ Osteoarchaeological Research Laboratory, University of Stockholm, Stockholm, Sweden.

¹⁰ Archaeological Institute, Research Centre for the Humanities, Hungarian Academy of Sciences, Budapest, Hungary.

¹¹ Zoological Institute RAS, Universitetskaya nab. 1, 199034 Saint-Petersburg, Russia

¹² The National Museum of Romanian History, 12 Calea Victoriei, 030026 Bucharest, Romania.

¹³ Institut de Génétique et Développement de Rennes, CNRS-UMR6290, Université de Rennes 1, Rennes, France.

¹⁴ Department of Archaeology, School of Geosciences, University of Aberdeen, St. Mary's, Elphinstone Road, AB24 3UF, UK.

* Corresponding authors: Laurent A. F. Frantz – laurent.frantz@arch.ox.ac.uk; Greger Larson – greger.larson@arch.ox.ac.uk; Dan G. Bradley – dbradley@tcd.ie

[†] Contributed equally

Abstract: The geographic and temporal origins of dogs remain controversial. Here, we generated genetic sequences from 59 ancient dogs and a complete (28x) genome of a late Neolithic dog (~4,800 calBP) from Ireland. Our analyses revealed a deep split separating modern East Asian and Western Eurasian dogs. Surprisingly, the date of this divergence (~14,000-6,400 years ago) occurs commensurate or several millennia after the first appearance of dogs in Europe and East Asia. Additional analyses of ancient and modern mitochondrial DNA revealed a sharp discontinuity in haplotype frequencies in Europe. Combined, these results suggest that dogs may have been domesticated independently in Eastern and Western Eurasia from distinct wolf populations. East Eurasia dogs were then possibly transported alongside people where they partially replaced European Palaeolithic dogs.

One Sentence Summary: Genomics and archaeology reveal both a possible dual origin of domestic dogs and a subsequent translocation of East Asian dogs into Europe.

Main Text: Dogs were the first domestic animal and the only animal domesticated prior to the advent of settled agriculture (1). Despite their importance in human history, no consensus has emerged with regard to their geographic and temporal origins, or whether dogs were domesticated just once or independently on more than one occasion. Though several claims have been made for an initial appearance of dogs in the early Upper Palaeolithic (~30,000 years ago; *e.g.* 2), the first remains confidently assigned to dogs appear in Europe ~15,000 years ago and in Far East Asia over 12,500 years ago (1, 3). While archaeologists remain open to the idea that there was more than one geographic origin for dogs (*e.g.* (4, 5), most genetic studies have concluded that dogs were likely domesticated just once (6) – disagreeing on whether this occurred in Europe (7), Central Asia (8), or East Asia (9).

Recent palaeogenetic studies have had a tremendous impact on our understanding of early human evolution (*e.g.* (10, 11)). Here we apply a similar approach to reconstruct the evolutionary history of dogs. We generated 59 ancient mtDNA sequences from European dogs (from 14,000 to 3,000 years ago) as well as a high coverage nuclear genome (~28x) of an ancient dog ~4,800 calBP (12) from the Neolithic passage grave complex of Newgrange (*Sí an Bhrú*) in Ireland. We combined our ancient sample with 80 modern publically available full genome sequences and 605 modern dogs (including village dogs and 48 breeds) genotyped on the 170k HD SNP array (12).

We first assessed characteristics of the Newgrange dog by typing SNPs associated with specific phenotypic traits and by inferring its level of inbreeding, compared to other breed and village dogs (12). Our results suggest that the degree of artificial selection and controlled breeding during the Neolithic was similar to that observed in modern free-living dogs. In addition, the Newgrange dog did not possess variants associated with modern breed-defining traits including hair length or coat color. And though this dog was likely able to digest starch less efficiently than modern dogs, it was more efficient than wolves (12).

A phylogenetic analysis, based on 170k SNPs revealed a deep split separating the modern Sarloos breed from other dogs (Fig. 1a). This breed - created in the 1930s in the Netherlands - involved breeding German Shepherds with captive wolves (13), thus explaining the breed's topological placement. Interestingly, the second deepest split (evident on the basis of both the 170K SNP panel – Fig 1a - and genome-wide SNPs - Fig. S4) separates modern East Asian and Western Eurasian (Europe and the Middle East) dogs. Moreover, the Newgrange dog clusters tightly with Western Eurasian dogs. We used Principal Component Analysis (PCA), D-statistics and *TreeMix* (12) to further test this pattern. Each of these analyses unequivocally placed the Newgrange dog with modern European dogs (Figs. S5, S6, S7). These findings demonstrate that the node separating the East Asian and Western Eurasian clades is older than the Newgrange individual; directly radiocarbon dated to ~4,800 years ago.

Other nodes leading to multiple dog populations and breeds (including the basal breeds (1) such as Greenland Sledge dogs or Siberian Husky; Fig. 1a) are poorly supported, suggesting that these breeds likely possess mixed ancestry from both Western Eurasian and East Asian dog lineages. To further assess the robustness of the deep split and those nodes associated with the potentially admixed lineages, we defined Western Eurasian and East Asian “core” groups (Fig. 1a) supported by the strength of the node leading to each cluster (12). We then used D-statistics to assess the affinity of each population to either Western Eurasian or East Asian core groups (12). The results of this analysis again revealed a clear East-West geographic pattern across Eurasia associated with the deep phylogenetic split (Fig. 1b). Breeds such as the Eurasier, Greenland Sledge dogs and Siberian Huskies (all basal breeds from Northern regions(1)), however, possess strong signatures of admixture with the East Asian core samples (Fig. S11), as do populations sampled in East Asia that clustered alongside Western Eurasian dogs (e.g. Papua New Guinean village dog; Fig. 1a).

We used the Multiple Sequentially Markovian Coalescent (*MSMC*)(12, 14) to reconstruct the population history of East Asian and Western Eurasia dogs. An analysis of individual high coverage genomes demonstrated a long, shared population history between the Newgrange dog and modern dogs from both Western Eurasia and East Asia (Fig. S15). A reconstruction using two genomes per group improved the resolution for recent time periods (Fig. 2a) and revealed a bottleneck in the Western Eurasian population, following its divergence from the East Asian core. A similar bottleneck observed in non-African human populations has been interpreted as a signature of a migration out of Africa (15). We therefore speculate that the analogous bottleneck observed in our dataset could be the result of a divergence and subsequent migration from east to west; supporting suggestions drawn from recent analyses of modern dog genomes (8, 9, 16).

We then used *MSMC* to compute divergence times as a mean to assess the time frame of the shared population history among dogs, and between dogs and wolves. To obtain reliable time estimates, we used the radiocarbon age of the Newgrange dog to calibrate the mutation rate for dogs (12)(Fig. S16). This resulted in a mutation rate estimate of between 0.3×10^{-8} and 0.45×10^{-8} per generation - similar to that obtained with an ancient grey wolf genome (17). Using this mutation rate, we calculated the divergence time between the two modern Russian wolves (18) used in this study and the modern dogs to be 60,000-20,000 years ago (Fig. S17; Fig. 2b). Importantly, this date should not be interpreted as a time frame for domestication, since the wolves we examined may not have been closely related to the population that gave rise to dogs (6).

124 These analyses also suggested that the divergence between the East Asian and Western
125 Eurasian core groups (~14,000-6,400 years ago) occurred commensurate, or several millennia
126 after the earliest known appearance of domestic dogs in both Europe (>15,000 years) and East
127 Asia (>12,500 years) (1) (Figs. S17, 2b). In addition, admixture signatures from wolves into
128 Western Eurasian dogs most likely pushed this estimated time of divergence deeper into the past
129 (12) meaning that the expected time of divergence between East and Western cores is likely
130 younger than our estimate. These results imply that indigenous populations of dogs were already
131 present in Europe and East Asia during the Palaeolithic (prior to this genomic divergence).
132 Under this hypothesis, this early indigenous dog population in Europe was replaced (at least
133 partially) by the arrival of East Eurasian dogs.

134 To investigate this potential replacement, we sequenced and analyzed 59 hyper-variable
135 mtDNA fragments from ancient dogs spread across Europe and combined those with 167
136 modern sequences (12). Each sequence was then assigned to one of four major well-supported
137 haplogroups (A-D) (19). While the majority of ancient European dogs belonged to either
138 haplogroup C or D (63% and 20%, respectively), most modern European dogs possess sequences
139 within haplogroups A and B (64 and 22% respectively) (Fig. 2c, d, e). Using simulations, we
140 showed that this finding cannot be explained by drift alone (12). Instead, this pattern arose from
141 clear turnover in the mitochondrial ancestry of European dogs, most likely as a result of an
142 arrival of East Asian dogs. This migration led to a partial replacement of ancient dog lineages in
143 Europe that were present by at least 15,000 years ago (1).

144 Though the mtDNA turnover is obvious, the nuclear signature reveals an apparent long-
145 term continuity. Assessments of ancestry in humans have demonstrated that major (nuclear)
146 turnovers can be difficult to detect without samples from the admixing population (11). A
147 genome-wide PCA analysis revealed that PC2 clearly discriminates the Newgrange dog from
148 other modern dogs (Fig. S8), suggesting that this individual possessed ancestry from an
149 unsampled population.

150 Our MSMC analysis reveals that the population split between the Newgrange dog and the
151 East Asian core (as measured by cross coalescence rate [CCR]) is older (on average) than the
152 split between modern Western Eurasian and East Asian lineages (Fig. 2b). Simulations suggest
153 that this pattern could be explained by a partial replacement model in which the Newgrange dog
154 retained a degree of ancestry from an outgroup population (Fig. S20a,b), that was different from
155 modern wolves (12). Alternatively, this pattern could also be explained by secondary gene flow
156 from Asian dogs into modern European dogs (Fig. S20c). Nevertheless, simulations show that
157 secondary gene flow has a smaller effect on CCR than the partial replacement model (Fig.
158 S20b,d). Moreover, secondary gene flow cannot explain the placement of the Newgrange dog on
159 our genome-wide PCA (Fig. S8). Overall, these observations are consistent with a scenario in
160 which the Newgrange dog retained a degree of ancestry from an ancient canid population that
161 falls outside of the variation of modern dogs, but that is also different from modern wolves. This
162 pattern also suggests that the replacement of European indigenous Palaeolithic dogs may not
163 have been complete.

164 To assess the consilience between our results and the archaeological record, we compiled
165 evidence for the earliest dog remains across Eurasia (Fig. 3a). We found that while dogs are
166 present at sites as old as 12,500 years in Eastern Eurasia (China, Kamchatka and East Siberia)
167 and 15,000 years in Western Eurasia (Europe and Near East) dog remains older than 8,000 years
168 have yet to be recovered in Central Eurasia (Fig. 3a; Table S7). Combined with our DNA

analyses, this observation suggests that two distinct populations of dogs were present in Eastern and Western Eurasia during the Palaeolithic.

The establishment of these populations is consistent with two scenarios: a single origin of Eurasian dogs followed by early transportation, founder effects, isolation and drift, or two independent domestication processes on either side of Eurasia. In the first scenario, the archaeological record should reveal a temporal cline of the first appearance of dogs across Eurasia stemming from a single source. Given the current lack of dog remains prior to 8,000 years ago in Central Eurasia, a scenario involving a single origin followed by an early transportation seems less likely.

Given our combined results, we suggest the following hypothesis: two genetically differentiated and potentially extinct wolf populations in Eastern (8, 9) and Western Eurasia (7) may have been independently domesticated prior to the advent of settled agriculture (Fig. 3a). The eastern dog population then dispersed westward alongside humans, between 6,400 and 14,000 years ago, into Western Europe (10, 11, 20) whereupon they partially replaced an indigenous Palaeolithic dog population. Our hypothesis reconciles previous studies that have suggested domestic dogs originated in East Asia (9, 19) and Europe (7). For numerous reasons, the null hypothesis should be that individual animal species were domesticated just once (21). The combined genetic and archaeological results presented here, however, suggest that dogs, like pigs (22), may have been domesticated twice. Nevertheless, given the complexity of the evolutionary history of dogs and uncertainties related to mutation rates, generation times and the incomplete nature of the archaeological record, our scenario remains hypothetical. Genome sequences derived from ancient Eurasian dogs and wolves will provide the necessary means to assess whether dog domestication occurred more than once.

References and Notes:

1. G. Larson *et al.*, *Proc. Natl. Acad. Sci.* **109**, 8878–83 (2012).
2. M. Germonpré, M. Lázníčková-Galetová, M. V. Sablin, *J. Archaeol. Sci.* **39**, 184–202 (2012).
3. M. Pionnier-Capitan *et al.*, *J. Archaeol. Sci.* **38**, 2123–2140 (2011).
4. T. Dayan, *J. Archaeol. Sci.* **21**, 633–640 (1994).
5. M. Ollivier *et al.*, *PLoS One*. **8**, e75110 (2013).
6. A. H. Freedman *et al.*, *PLoS Genet.* **10**, e1004016 (2014).
7. O. Thalmann *et al.*, *Science*. **342**, 871–4 (2013).
8. L. M. Shannon *et al.*, *Proc. Natl. Acad. Sci. U. S. A.* **112**, 13639–13644 (2015).
9. G.-D. Wang *et al.*, *Cell Res.* **26**, 21–33 (2015).
10. M. E. Allentoft *et al.*, *Nature*. **522**, 167–172 (2015).
11. W. Haak *et al.*, *Nature*. **522**, 207–211 (2015).
12. Supplementary Material.
13. D. Morris, *Dogs: The Ultimate Dictionary of Over 1,000 Dog Breeds* (Trafalgar Square, 2008).

- 208 14. S. Schiffels, R. Durbin, *Nat. Genet.* **46**, 919–25 (2014).
- 209 15. H. Li, R. Durbin, *Nature*. **475**, 493–6 (2011).
- 210 16. M. Pilot *et al.*, *Proc. Biol. Sci.* **282**, 20152189– (2015).
- 211 17. P. Skoglund, E. Ersmark, E. Palkopoulou, L. Dalén, *Curr. Biol.* **25**, 1515–1519 (2015).
- 212 18. G. Wang *et al.*, *Nat. Commun.* **4**, 1860 (2013).
- 213 19. P. Savolainen, Y. Zhang, J. Luo, J. Lundeberg, T. Leitner, *Science*. **298**, 1610–3 (2002).
- 214 20. L. M. Cassidy *et al.*, *Proc. Natl. Acad. Sci.* **113**, 201518445 (2015).
- 215 21. G. Larson, J. Burger, *Trends Genet.* **29**, 197–205 (2013).
- 216 22. L. A. F. Frantz *et al.*, *Nat. Genet.* **47**, 1141–1148 (2015).
- 217 23. C. Gamba *et al.*, *Nat. Commun.* **5**, 5257 (2014).
- 218 24. L. H. van Wijngaarden-Bakker, *Proc. R. Irish Acad. Sect. C Archaeol. Celt. Stud. Hist.*
219 *Linguist. Lit.* **86C**, 17–111 (1986).
- 220 25. M. J. O’Kelly, R. M. Cleary, D. Lehané, *Newgrange, Co. Meath, Ireland: The Late*
221 *Neolithic/Beaker Period Settlement* (B.A.R., 1983).
- 222 26. F. McCormick, in *The Holocene History of the European Vertebrate Fauna: Modern*
223 *Aspects of Research*, N. Benecke, Ed. (1998).
- 224 27. P. Reimer, *Radiocarbon*. **55**, 1869–1887 (2013).
- 225 28. S. Pääbo *et al.*, *Annu. Rev. Genet.* **38**, 645–679 (2004).
- 226 29. L. Orlando *et al.*, *Genome Res.* **21**, 1705–19 (2011).
- 227 30. J. T. Vilstrup *et al.*, *PLoS One*. **8**, e55950 (2013).
- 228 31. M. Meyer, M. Kircher, *Cold Spring Harb. Protoc.* **2010**, pdb.prot5448 (2010).
- 229 32. M. Martin, *EMBnet.journal*. **17**, 10 (2011).
- 230 33. H. Li, R. Durbin, *Bioinformatics*. **25**, 1754–60 (2009).
- 231 34. K. Lindblad-Toh *et al.*, *Nature*. **438**, 803–19 (2005).
- 232 35. M. Schubert *et al.*, *Nat. Protoc.* **9**, 1056–82 (2014).
- 233 36. H. Li *et al.*, *Bioinformatics*. **25**, 2078-9 (2009).
- 234 37. A. McKenna *et al.*, *Genome Res.* **20**, 1297–1303 (2010).
- 235 38. K. Okonechnikov, A. Conesa, F. García-Alcalde, *Bioinformatics*. **32**, 292-4 (2015).
- 236 39. H. Jónsson, A. Ginolhac, M. Schubert, P. L. F. Johnson, L. Orlando, *Bioinformatics*. **29**,
237 1682–4 (2013).
- 238 40. B. Bai *et al.*, *Nucleic Acids Res.* **43**, 777-83 (2014).
- 239 41. A. Auton *et al.*, *Science*. **336**, 193-8 (2012).
- 240 42. A. R. Quinlan, I. M. Hall, *Bioinformatics*. **26**, 841–2 (2010).
- 241 43. E. Han, J. S. Sinsheimer, J. Novembre, *Mol. Biol. Evol.* **31**, 723–35 (2014).

242 44. A. Vaysse *et al.*, *PLoS Genet.* **7**, e1002316 (2011).

243 45. M. L. Speir *et al.*, *Nucleic Acids Res.*, **44** 717–25 (2016).

244 46. S. Purcell *et al.*, *Am. J. Hum. Genet.* **81**, 559–75 (2007).

245 47. E. Paradis, J. Claude, K. Strimmer, *Bioinformatics.* **20**, 289–290 (2004).

246 48. N. Patterson, A. L. Price, D. Reich, *PLoS Genet.* **2**, e190 (2006).

247 49. J. K. Pickrell, J. K. Pritchard, *PLoS Genet.* **8**, e1002967 (2012).

248 50. E. Y. Durand, N. Patterson, D. Reich, M. Slatkin, *Mol. Biol. Evol.* **28**, 2239–2252 (2011).

249 51. N. Patterson *et al.*, *Genetics.* **192**, 1065–93 (2012).

250 52. S. R. Browning, B. L. Browning, *Am. J. Hum. Genet.* **81**, 1084–97 (2007).

251 53. Z. Fan *et al.*, *Genome Res.* **26**, 163–73 (2016).

252 54. H. Angleby, P. Savolainen, *Forensic Sci. Int.* **154**, 99–110 (2005).

253 55. A. Ardalan *et al.*, *Ecol. Evol.* **1**, 373–85 (2011).

254 56. A. R. Boyko *et al.*, *Proc. Natl. Acad. Sci. U. S. A.* **106**, 13903–8 (2009).

255 57. S. K. Brown *et al.*, *PLoS One.* **6**, e28496 (2011).

256 58. S. Castroviejo-Fisher, P. Skoglund, R. Valadez, C. Vilà, J. A. Leonard, *BMC Evol. Biol.*
257 **11**, 73 (2011).

258 59. V. Muñoz-Fuentes, C. T. Darimont, P. C. Paquet, J. A. Leonard, *Conserv. Genet.* **11**, 547–
259 556 (2009).

260 60. N. Okumura, N. Ishiguro, M. Nakano, A. Matsui, M. Sahara, *Anim. Genet.* **27**, 397–405
261 (2009).

262 61. M. C. R. Oskarsson *et al.*, *Proc. Biol. Sci.* **279**, 967–74 (2012).

263 62. J.-F. Pang *et al.*, *Mol. Biol. Evol.* **26**, 2849–64 (2009).

264 63. P. Savolainen, T. Leitner, A. N. Wilton, E. Matisoo-Smith, J. Lundeberg, *Proc. Natl.*
265 *Acad. Sci. U. S. A.* **101**, 12387–90 (2004).

266 64. K. Tsuda, Y. Kikkawa, H. Yonekawa, Y. Tanabe, *Genes Genet. Syst.* **72**, 229–38 (1997).

267 65. J. A. Leonard *et al.*, *Science.* **298**, 1613–6 (2002).

268 66. M. F. Deguilloux, J. Moquel, M. H. Pemonge, G. Colombeau, *J. Archaeol. Sci.* **36**, 513–
269 519 (2009).

270 67. F. Verginelli *et al.*, *Mol. Biol. Evol.* **22**, 2541–51 (2005).

271 68. M. Pionnier-Capitan, thesis, ENS Lyon (2010).

272 69. O. Lebrasseur, thesis, Durham University (2014).

273 70. N. Rohland, M. Hofreiter, *Nat. Protoc.* **2**, 1756–62 (2007).

274 71. A. Cooper, *Science (80-.)*. **289**, 1139b–1139 (2000).

275 72. M. T. P. Gilbert, H.-J. Bandelt, M. Hofreiter, I. Barnes, *Trends Ecol. Evol.* **20**, 541–4

- (2005).
73. K. S. Kim, S. E. Lee, H. W. Jeong, J. H. Ha, *Mol. Phylogenet. Evol.* **10**, 210–20 (1998).
74. M.-S. Peng *et al.*, *Mol. Ecol. Resour.* **15**, 1238–42 (2015).
75. R. C. Edgar, *Nucleic Acids Res.* **32**, 1792–7 (2004).
76. J. Leigh, PopART (Population Analysis with Reticulate Trees) (2015).
77. R. R. Hudson, *Bioinformatics.* **18**, 337–338 (2002).
78. G. K. Chen, P. Marjoram, J. D. Wall, *Genome Res.* **19**, 136–42 (2009).
79. D. H. Alexander, J. Novembre, K. Lange, *Genome Res.* **19**, 1655–64 (2009).
80. M. Boudadi-Maligne, J.-B. Mallye, M. Langlais, C. Barshay-Szmidt, *PALEO*, 39–54 (2012).
81. R. J. Losey *et al.*, *PLoS One.* **8**, e63740 (2013).
82. N. Benecke, *J. Archaeol. Sci.* **14**, 31–49 (1987).
83. M. Germonpré *et al.*, *J. Archaeol. Sci.* **36**, 473–490 (2009).
84. R. Flad, *J. East Asian Archaeol.* **3**, 23–51 (2001).
85. L. Barton *et al.*, *Proc. Natl. Acad. Sci. U. S. A.* **106**, 5523–8 (2009).
86. A. Lasota-Moskalewska, K. Szymczak, M. Khudzhazarov, *Archaeol. Balt.* **11** (2009).
87. A. Razzokov. *Sarazm. Dushanbe: Institute of History, Archaeology, and Ethnography. Dushanbe, Tajikistan: Academy of Sciences of Tajikistan A. Donish History, Archaeology and Ethnographic Institute.* (2008).
88. H. Bocherens, M. Mashkour, D. Billiou, *Environ. Archaeol.* **33**, 253–64 (2013).
89. A. K. Kasparov, *Paléorient.* **22**, 161–167 (1996).
90. M. Frachetti, N. Benecke, *Antiquity.* **83**, 1023–1037 (2015).
91. C. Chang, N. Benecke, F. Grigoriev, A. Rosen, P. Tourtellotte, *Antiquity.* **77**, 298–312 (2003).
92. V. Bakhshaliev, in *The Archaeology of Nakhichevan. Ten Years of New Discoveries* (Ege Yayinlari, Istanbul, 1997).
93. R. Berthon *et al.*, *Environ. Archaeol.* **18**, 191–200 (2013).
94. S. J. Pawankar and P. K. Thomas, *Bulletin of the Deccan College Research Institute* **56**, 363–370 (1996).
95. B. & R. Allchin, F. R. Allchin, in *The Birth of Indian Civilization: India and Pakistan Before 500 B.C.* (1968).
96. R. V. Joshi, *Archaeol. Surv. India, New Delhi, India*, 12–16 (1961).
97. P. K. Thomas, in *The Walking larder: patterns of domestication, pastoralism, and predation* (Routledge, 2014).
98. B. P. Sahu, *From hunters to breeders: faunal background of early India* (Anamika

- Prakashan, 1988).
99. N. Bhola, G. V. S. Rao, *Animal remains from Lothal excavations* (Zoological Survey of India, 1962).
 100. R. B. . Sewell, B. S. Guha, in *Mohenjo-Daro and the Indus* (1931).
 101. G. Stein, in *Early Animal Domestication and Its Cultural Context*, P. J. Crabtree, D. V. Campana, K. Ryan, Eds. (UPenn Museum of Archaeology, 1989).
 102. Telegin D.Ya., *redno-Stogivska kultura epokhi midi*. (1973).
 103. K. N. Wilkinson *et al.*, *J. F. Archaeol.* **37**, 20-33 (2013).
 104. R. D. Barnett, W. Watson, *Iraq.* **14**, 132 (1952).
 105. M. D. Frachetti, N. Benecke, A. N. Mar'yashev, P. N. Doumani, *World Archaeol.* **42**, 622–646 (2010).
 106. P. M. Dolukhanov, in *Hunters in transition: Mesolithic societies of temperate Eurasia and their transition to farming* (1986).
 107. G. Matyushin, *The mesolithic and neolithic in the southern Urals and central Asia*. (1986).
 108. N. D. Ovodov *et al.*, *PLoS One.* **6**, e22821 (2011).
 109. M. Germonpré, M. Lázníková-Galetová, R. J. Losey, J. Rääkkönen, M. V. Sablin, *Quat. Int.* **359-360**, 261–279 (2015).
 110. M.-A. Garcia, *Bull. la Société préhistorique française.* **102**, 103–108 (2005).
 111. M. V. Sablin, G. A. Khlopachev, *Curr. Anthropol.* **43**, 795–799 (2002).
 112. A. Perri, *J. Archaeol. Sci.* **68**, 1–4 (2016).
 113. S. J. Crockford, Y. V. Kuzmin, *J. Archaeol. Sci.* **39**, 2797–2801 (2012).
 114. M. Boudadi-Maligne, G. Escarguel, *J. Archaeol. Sci.* **45**, 80–89 (2014).
 115. A. G. Drake, M. Coquerelle, G. Colombeau, *Sci. Rep.* **5**, 8299 (2015).
 116. D. P. Howrigan, M. A. Simonson, M. C. Keller, *BMC Genomics.* **12**, 460 (2011).
 117. E. Axelsson *et al.*, *Nature.* **495**, 360–4 (2013).
 118. E. Cadieu *et al.*, *Science.* **326**, 150–3 (2009).
 119. H. G. Parker *et al.*, *Science.* **325**, 995–8 (2009).

Acknowledgments: Raw reads of the Newgrange dog have been deposited at the European Nucleotide Archive (ENA) with project number: PRJEB13070. Mitochondrial sequences as well as genotype files (in plink format) were deposited on Dryad (<http://datadryad.org/review?doi=doi:10.5061/dryad.8gp06>). We would like to thank G. Wang, J. Schraiber, L. Orlando, L. Dalén, R.E. Green, P. Savolainen, E. Loftus for their valuable input. We are also grateful to A. Oszrás and I. Zalai-Gaál R.-M. Arbogast, A. Beeching, A. Boroneant, O. Lecomte, S. Madeleine, C. & D. Mordant, M. Patou-Mathis, P. Pétrequin, L. Salanova, J. Schibler, A. Tsuneki, F. Valla for providing archaeological material. L.A.F.F., O.L., A.L. and

G.L. were supported by a European Research Council grant (ERC-2013-StG-337574-UNDEAD) and Natural Environmental Research Council grants (NE/K005243/1 and NE/K003259/1). L.A.F.F. was supported by a Junior Research Fellowship (Wolfson College, University of Oxford). V.E.M, V.M , M.D.T. and the sequencing of the Newgrange dog genome were funded by ERC Investigator grant 295729-CodeX awarded to D.G.B. We would also like to acknowledge the National Museum of Ireland for providing the petrous bone of the Newgrange dog and the Science Foundation Ireland Award 12/ERC/B2227 and Trinseq. AB was supported by a Romanian National Authority for Scientific Research (PN-II-ID-PCE-2011-3-1015). The work at ENS Lyon and at MNHN Paris was also supported by PURINA-NESTLE. The authors declare no conflict of interest.

Fig. 1: Deep split between East Asian and Western Eurasian dogs. *a.* A neighbour-joining tree (with bootstrap values) based on Identity by State (*I2*) of 605 dogs. Red and yellow clades represent the East Asian and Western Asian core groups respectively (*I2*). *b.* A map showing the location and relative proportion of ancestry (mean *D*-values) of dogs (Fig. S10). Positive values (red) indicate that the population shares more derived alleles with the East Asian core while negative values (yellow) indicate a closer association with the Western Eurasian core.

Fig. 2: Effective population size, divergence times and mtDNA. *a.* Effective population size through time of East and Western Eurasian dogs and wolves with MSMC. *b.* Cross-coalescence rate (CCR) per year for each population pair in Fig. 2a. The CCR represents the ratio of within and between population coalescence rates (CR). The ratio measures the age and pace of divergence between two populations. Values close to 1 indicate that both within and between CR are equal meaning the two populations have not yet diverged. Values close to 0 indicate that the populations have completely diverged. *c.* Bar plot representing the proportion of mtDNA haplogroups at different time periods. *d.* Locations of archaeological sites with haplogroup proportions. *e.* Location of modern samples with haplogroup proportions.

Fig. 3: Archaeological evidence for the first appearance of dogs across Eurasia and a model of dog domestication. *a.* Map representing the geographic origin and age of the oldest archaeological dog remains in Eurasia (*I2*). *b.* A suggested model of dog domestication under the dual origin hypothesis. An initial wolf population split into East and West Eurasian wolves that were then domesticated independently before going extinct (as indicated by the † symbol). The Western Eurasian dog population (European) was then partially replaced by a human-mediated translocation of Asian dogs at least 6,400 years ago, a process that took place gradually after the arrival of the eastern dog population.

Supplementary Materials:

Materials and Methods

Figs. S1-S29

Tables S1-S7

References (23-110)